



**9th Conference of the
Society of Physicists of Macedonia**

BOOK OF ABSTRACTS

20 - 23 September 2012
Ohrid, Macedonia

Conference Chair

Blagoja Veljanoski (Macedonia)

Scientific Committee

President	Olga Galbova (Macedonia)
Members	Bayram Akarsy (Turkey)
	Dan Cojoc (Italy)
	Elena Atanasova (Bulgaria)
	Gertrud Zwicknagl (Germany)
	Josip Slisko Ignjatov (Mexico)
	Ljupco Hadzievski (Serbia)
	Milko Jaksic (Croatia)
	Mimoza Ristova (Macedonia)
	Nenad Novkovski (Macedonia)
	Ninoslav Stojadinovic (Serbia)
	Oliver Zajkov (Macedonia)
	Valentin Peschansky (Ukraine)

Organizing Committee

President	Stojan Manolev (Macedonia)
Members	Irina Petreska (Macedonia)
	Lihnida Stojanovska Georgievska (Macedonia)
	Suzana Topuzoski (Macedonia)
	Olgica Kuzmanovska Barandovska (Macedonia)
	Aleksandar Skeparovski (Macedonia)
	Ane Tuntev (Macedonia)
	Fadil Ajredini (Macedonia)
	Mirjana Davkova (Macedonia)

Technical Support

Jovan Stanoevski
Vesna Manchevska
Stefan Terpini

**EVALUATION OF POPULATION DOSES IN
STATISTICAL REGIONS OF MACEDONIA
RESULTING FROM THE NATURAL
RADIOACTIVITY IN THE SOIL**

Zdenka Stojanovska¹, Zora S. Zunic², Mimoza Ristova³

¹Faculty of Medical Sciences, Goce Delcev University, Stip,
Republic of Macedonia

²Institute of Nuclear Sciences "Vinca", University of Belgrade,
Serbia

³Institute of Physics, Faculty of Natural Sciences and
Mathematic, University in Skopje, Republic of Macedonia

Terrestrial radiation from naturally occurring radionuclides, such as ^{40}K , radionuclides from the ^{238}U and ^{232}Th series and their decay products is the main external source of public exposure. This research presents the results of assessment the population doses from natural radionuclides in soils at the eight statistical regions of the Republic of Macedonia. The surface soil samples (from 0 to 20 cm depth) were collected during the period 2007-2010 from 213 locations of the entire country territory. High-resolution gamma spectroscopy was employed the qualitative and quantitative analysis. The obtained specific activity of ^{40}K , ^{226}Ra , ^{232}Th and associated external dose resulting from the gamma-radiation were different in the soil of each region depending of the geological and geographical conditions. The results showed that the obtained values for gamma absorbed dose rate in air varied from 50 nGy h⁻¹ in Skopje to 86 nGy h⁻¹ in Pelagonija. The arithmetic and geometric mean values of the annual effective dose equivalent for the Republic of Macedonia were found to be: 83 $\mu\text{Sv y}^{-1}$ (SD=29) and 79 $\mu\text{Sv y}^{-1}$ (GSD=1.4), respectively.